Table 2.10C Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Numbers in Thousands, 2010 and 2011

Dependence or Abuse	Total ¹ (2010)	Total ¹ (2011)	MDE (2010)	MDE (2011)	No MDE (2010)	No MDE (2011)
DEPENDENCE ²	(2010)	(===)	(2010)	(===1)	(2010)	(====)
Illicit Drugs ³	34	37	15	17	30	32
Marijuana	29	32	12	13	26	28
Illicit Drugs Other Than Marijuana ³	22	20	11	13	17	15
Alcohol	30	26	14	12	26	23
Both Illicit Drugs and Alcohol ³	16	14	8	9	14	11
Illicit Drugs or Alcohol ³	42	43	18	19	38	39
DEPENDENCE OR ABUSE ²						
Illicit Drugs ³	47	47	21	21	42	43
Marijuana	40	41	15	16	36	38
Illicit Drugs Other Than Marijuana ³	30	29	18	16	24	24
Alcohol	49	45	20	21	45	40
Both Illicit Drugs and Alcohol ³	31	30	15	14	26	26
Illicit Drugs or Alcohol ³	60	58	25	27	57	54

^{*}Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Table 2.10D Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2010 and 2011

	Total ¹	Total ¹	MDE	MDE	No MDE	No MDE
Dependence or Abuse	(2010)	(2011)	(2010)	(2011)	(2010)	(2011)
DEPENDENCE ²						
Illicit Drugs ³	0.14	0.15	0.77	0.81	0.13	0.14
Marijuana	0.12	0.13	0.60	0.63	0.12	0.12
Illicit Drugs Other Than Marijuana ³	0.09	0.08	0.58	0.65	0.08	0.07
Alcohol	0.12	0.10	0.69	0.57	0.12	0.10
Both Illicit Drugs and Alcohol ³	0.07	0.06	0.41	0.43	0.06	0.05
Illicit Drugs or Alcohol ³	0.17	0.17	0.93	0.91	0.17	0.17
DEPENDENCE OR ABUSE ²						
Illicit Drugs ³	0.19	0.19	1.05	0.97	0.18	0.19
Marijuana	0.16	0.16	0.79	0.76	0.16	0.17
Illicit Drugs Other Than Marijuana ³	0.12	0.12	0.92	0.78	0.11	0.11
Alcohol	0.20	0.18	1.00	1.00	0.20	0.17
Both Illicit Drugs and Alcohol ³	0.13	0.12	0.78	0.69	0.12	0.11
Illicit Drugs or Alcohol ³	0.25	0.23	1.18	1.21	0.24	0.23

^{*}Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Table 2.10P Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): P Values from Tests of Differences of Percentages, 2011 Versus 2010

	Total ¹	Total ¹	MDE	MDE	No MDE	No MDE
Dependence or Abuse	(2010)	(2011)	(2010)	(2011)	(2010)	(2011)
DEPENDENCE ²						
Illicit Drugs ³	0.9324		0.7812		0.9324	
Marijuana	0.9286		0.7542		0.7552	
Illicit Drugs Other Than Marijuana ³	0.4113		0.9697		0.3200	
Alcohol	0.1429		0.1800		0.3566	
Both Illicit Drugs and Alcohol ³	0.2601		0.6374		0.3069	
Illicit Drugs or Alcohol ³	0.5433		0.3514		0.8611	
DEPENDENCE OR ABUSE ²						
Illicit Drugs ³	0.6955		0.1824		0.7071	
Marijuana	0.8586		0.8488		0.9084	
Illicit Drugs Other Than Marijuana ³	0.2850		0.2529		0.6721	
Alcohol	0.0050		0.4261		0.0098	
Both Illicit Drugs and Alcohol ³	0.0210		0.1623		0.0775	
Illicit Drugs or Alcohol ³	0.1793		0.3505		0.3951	

^{*}Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.